

Quasigroups, Asymptotic Symmetries and Conservation Laws in General Relativity

Alexander I. Nesterov

Departament of Physics, C.U.C.E.I., Guadalajara University, Guadalajara, Jalisco, Mexico*

(Dated: February 7, 2008)

A new quasigroup approach to conservation laws in general relativity is applied to study asymptotically flat at future null infinity spacetime. The infinite-parametric Newman-Unti group of asymptotic symmetries is reduced to the Poincaré quasigroup and the Noether charge associated with any element of the Poincaré quasialgebra is defined. The integral conserved quantities of energy-momentum and angular momentum are linear on generators of Poincaré quasigroup, free of the supertranslation ambiguity, possess the flux and identically equal to zero in Minkowski spacetime.

PACS numbers: 04.20.Ha, 04.20.-q

Keywords: nonassociativity, Poincaré quasigroup, asymptotic symmetries, conservation laws

As well known the general covariance of the Einstein equations results in differential conservation laws related to the field equations. From Noether's theorem one shows that for an arbitrary diffeomorphism generated by the vector field ξ the invariance of the Lagrangian for the Einstein equations leads to the conservation laws of the form

$$\mathfrak{J}^\mu(\xi)_{;\mu} = 0, \quad (1)$$

where the vector density $\mathfrak{J}^\mu$ is defined as $\mathfrak{J}^\mu(\xi) = \mathfrak{h}^{\mu\nu}_{;\nu}$, $\mathfrak{h}^{\mu\nu} = -\mathfrak{h}^{\nu\mu}$ being the *superpotential*, which is constructed from the densities of spin, bispin, vector field ξ and its derivatives [1, 2, 3]. For the Einstein-Hilbert action one obtains up to a factor of 2 a simple expression

$$\mathfrak{J}^\mu = -\frac{1}{4\pi} \left(\sqrt{-g} \xi^{[\mu;\nu]} \right)_{;\nu} \quad (2)$$

which was first given by Komar [4]. It is impossible simply to “renormalize” $\mathfrak{J}^\mu$ by factor of 2 because the normalization for energy-momentum and angular momentum differ by a factor of 2. The resolution of this problem is known [5]. One needs to add to the Einstein-Hilbert action I a surface term I_s and apply Noether's theorem to $I + I_s$. Komar's expression provides a fully satisfactory notion of the total mass in stationary, asymptotically flat spacetimes.

For an asymptotically flat spacetime the group of asymptotic symmetries is an infinite-parametric one. It contains an unique four-parametric translation subgroup and infinite-parametric subgroup of supertranslations. Both groups are the normal subgroups and the Lorentz group occurs as a factor group of the asymptotic symmetries group by the infinite dimensional subgroup of supertranslations. *Therefore there does not exist a canonical way of choosing the Poincaré group as a subgroup of the group of asymptotic symmetries. There are too many Poincaré subgroups, one for each supertranslations*

which is not translation. These circumstances generate the main difficulties in the numerous attempts to find the correct definition of angular momentum.

In our article a new quasigroup approach to the conservation laws developed in [2, 6, 7, 8, 9] is applied to asymptotically flat spacetime. The Poincaré quasigroup at future null infinity ($\mathcal{J}^+$) is introduced and compared with other definitions of asymptotic symmetries that have appeared in the literature. We define the Noether charge associated with any element of the Poincaré quasialgebra. It may be regarded as a form of linkages by Tamburino and Winicour [10] but with the new gauge conditions for asymptotic symmetries.

Quasigroups of transformations. The definition of the quasigroup of transformations first was given by Batalin [11]. Below we outline the main facts from the theory of the smooth quasigroups of transformations [12].

Let $\mathfrak{M}$ be a n -dimensional manifold and the continuous law of transformation is given by $x' = T_a x$, $x \in \mathfrak{M}$, where $\{a^i\}$ is the set of real parameters, $i = 1, 2, \dots, r$. The set of transformations $\{T_a\}$ forms a r -parametric quasigroup of transformations (with right action on $\mathfrak{M}$), if:

- 1) there exists a unit element which is common for all x^α and corresponds to $a^i = 0$: $T_a x|_{a=0} = x$;
- 2) the modified composition law holds:

$$T_a T_b x = T_{\varphi(b,a;x)} x;$$

- 3) the left and right units coincide:

$$\varphi(a, 0; x) = a, \quad \varphi(0, b; x) = b;$$

- 4) the modified law of associativity is satisfied:

$$\varphi(\varphi(a, b; x), c; x) = \varphi(a, \varphi(b, c; T_a x); x);$$

- 4) the transformation inverse to T_a exists: $x = T_a^{-1} x'$.
The generators of infinitesimal transformations

$$\Gamma_i = (\partial(T_a x)^\alpha / \partial a^i)|_{a=0} \partial / \partial x^\alpha \equiv R_i^\alpha \partial / \partial x^\alpha$$

form *quasialgebra* and obey the commutation relations

$$[\Gamma_i, \Gamma_j] = C_{ij}^p(x) \Gamma_p, \quad (3)$$

*Electronic address: nesterov@cencar.udg.mx

where $C_{ij}^p(x)$ are the structure functions satisfying the modified Jacobi identity

$$C_{ij,\alpha}^p R_k^\alpha + C_{jk,\alpha}^p R_i^\alpha + C_{ki,\alpha}^p R_j^\alpha + C_{ij}^l C_{kl}^p + C_{jk}^l C_{il}^p + C_{ki}^l C_{jl}^p = 0. \quad (4)$$

Theorem. Let the given functions R_i^α , C_{kj}^p obey the equations (3), (4), then locally the quasigroup of transformations is reconstructed as the solution of set of differential equations:

$$\frac{\partial \tilde{x}^\alpha}{\partial a^i} = R_j^\alpha(\tilde{x}) \lambda_i^j(a; x), \quad \tilde{x}^\alpha(0) = x^\alpha, \quad (5)$$

$$\frac{\partial \lambda_j^i}{\partial a^p} - \frac{\partial \lambda_p^i}{\partial a^j} + C_{mn}^i(\tilde{x}) \lambda_p^m \lambda_j^n = 0, \quad \lambda_j^i(0; x) = \delta_j^i. \quad (6)$$

Eq. (5) is an analog of the Lie equation, and Eq. (6) is the generalized Maurer-Cartan equation.

The notion of quasigroups is not widely known, therefore we give here two elementary examples playing the important role in the description of generalized coherent states for the groups $SU(2)$, $SU(1,1)$ and Thomas precession [16, 17].

Quasigroup $QS(2)$. This quasigroup is associated with the group $SU(2)$ and its action on the two-sphere S^2 . The sphere S^2 admits a natural quasigroup structure, namely, S^2 is a local two-parametric loop $QS(2)$ which is defined as follows. Let $\mathbb{C}$ be a complex plane and the isomorphism between points of the sphere and the complex plane $\mathbb{C}$ is established by the stereographic projection from the north pole of the sphere: $\zeta = e^{i\varphi} \tan(\theta/2)$. The nonassociative multiplication $\star$ is defined by

$$\zeta \star \eta = \frac{\zeta + \eta}{1 - \bar{\zeta}\eta}, \quad \zeta, \eta \in \mathbb{C}.$$

where a bar denotes complex conjugation.

Loop $QH(2)$. This loop is associated with the group $SU(1,1)$ and its action on the two-dimensional unit hyperboloid H^2 . Let $D \subset \mathbb{C}$ be the open unit disk, $D = \{\zeta \in \mathbb{C} : |\zeta| < 1\}$. We define the binary operation $*$ as following:

$$\zeta * \eta = \frac{\zeta + \eta}{1 + \bar{\zeta}\eta}, \quad \zeta, \eta \in D.$$

Inside D the set of complex numbers with the operation $*$ forms two-sided loop $QH(2)$, which is isomorphic to the geodesic loop of two-dimensional Lobachevskii space realized as the upper part of two-sheeted unit hyperboloid. The isomorphism is established by $\zeta = e^{i\varphi} \tanh(\theta/2)$, where (θ, φ) are inner coordinates on H^2 .

Null infinity and Poincaré quasigroup. It is known that for asymptotically flat at future null infinity ($\mathcal{J}^+$) space-time the group of asymptotic symmetries is the infinite-parametric Newman-Unti (NU) group which contains the infinite-dimensional Bondi-Metzner-Sachs (BMS) group preserving strong conformal geometry of $\mathcal{J}^+$ [18, 19, 20, 21].

NU group is the group of transformations $\mathcal{J}^+ \rightarrow \mathcal{J}^+$:

$$u \rightarrow u' = f(u, \zeta, \bar{\zeta}), \quad \partial f / \partial u > 0, \\ \zeta \rightarrow \zeta' = (\alpha\zeta + \beta) / (\gamma\zeta + \delta), \quad \alpha\delta - \beta\gamma = 1,$$

where $f(u, \zeta, \bar{\zeta})$ is an arbitrary function, ζ being a complex stereographic coordinates at two-dimensional space-like cross sections of $\mathcal{J}^+$ labeled by a coordinate u and its metric is given by

$$ds^2 = \frac{2d\zeta d\bar{\zeta}}{|P(u, \zeta, \bar{\zeta})|^2},$$

where we assume $P = VP_0$, $P_0 = (1/\sqrt{2})(1 + \zeta\bar{\zeta})$. When $V = 1$ this metric is reduced to the Bondi metric.

The infinite-parameter BMS-subgroup of NU-group is determined as following

$$u \rightarrow u' = K(\zeta, \bar{\zeta})(u + a(\zeta, \bar{\zeta})), \\ \zeta \rightarrow \zeta' = (\alpha\zeta + \beta) / (\gamma\zeta + \delta), \quad \alpha\delta - \beta\gamma = 1,$$

where $K(\zeta, \bar{\zeta}) = (1 + \zeta\bar{\zeta}) / (|\alpha\zeta + \beta|^2 + |\gamma\zeta + \delta|^2)$ and $a(\zeta, \bar{\zeta})$ is an arbitrary regular function on S^2 . The infinite-parameter normal subgroup of BMS-group

$$\zeta' = \zeta, \quad u' = u + a(\zeta, \bar{\zeta}),$$

is called the subgroup of *supertranslations* and contains a four-parameter normal translation subgroup

$$a = \frac{A + B\zeta + \bar{B}\bar{\zeta} + C\zeta\bar{\zeta}}{1 + \zeta\bar{\zeta}}.$$

The factor group of the BMS-group by the supertranslations consists from the conformal transformations $S^2 \rightarrow S^2$ and it is isomorphic to the *proper orthochronous Lorentz group*.

On $\mathcal{J}^+$ a general element of NU-algebra is given by (We use the spin coefficients formalism by Newman and Penrose [22] choosing $\kappa = \varepsilon + \bar{\varepsilon} = 0$, $\tau = \bar{\pi} = \bar{\alpha} + \beta$, $\rho = \bar{\rho}$)

$$\xi = B(u, \zeta, \bar{\zeta})\Delta^0 + C(u, \zeta, \bar{\zeta})\bar{\delta}^0 + \bar{C}(u, \zeta, \bar{\zeta})\delta^0, \quad (7)$$

where $\bar{\delta}C = 0$ and $\bar{\delta}, \Delta^0, \delta^0$ are the standard NP operators “edth”, Δ and δ restricted on $\mathcal{J}^+$.

In an arbitrary coordinate system on $\mathcal{J}^+$ the generators of four-parameter translation subgroup is given by

$$\xi_a = B_a(\zeta, \bar{\zeta})\Delta^0, \quad (C = 0), \quad (8)$$

where the function B_a is the solution of the following equation

$$\bar{\delta}^2 B_a = B_a(4(\bar{\alpha}^0)^2 - 2\bar{\delta}\bar{\alpha}^0),$$

(a runs from 1 to 4). The generators of “Lorentz group” is determined as following:

$$\xi_A = B_A(u, \zeta, \bar{\zeta})\Delta^0 + C_A\bar{\delta}^0 + \bar{C}_A\delta^0, \quad (\bar{\delta}C_A = 0), \quad (9)$$

where the function B_A satisfies (A runs from 1 to 6)

$$\dot{B}_A = Q_A(u, \zeta, \bar{\zeta}) + (1/2)(\bar{\partial}C_A + \partial\bar{C}_A),$$

here $Q_A(u, \zeta, \bar{\zeta})$ is an arbitrary real function and a dot being derivative with respect to the retarded time u . The generators of the NU-group obey the commutation relations:

$$\begin{aligned} [\xi_a, \xi_b] &= 0, & [\xi_a, \xi_B] &= C_{aB}^b(u, \zeta, \bar{\zeta})\xi_b, \\ [\xi_A, \xi_B] &= C_{AB}^D(u, \zeta, \bar{\zeta})\xi_D, \end{aligned} \quad (10)$$

where C_{aB}^b , C_{AB}^D are the *structure functions*, depending on arbitrary function Q [23].

Reduction of NU-group to the Poincaré quasigroup. The point is to obtain conditions on an arbitrary function B_A in the definition (9) of the generators of “Lorentz group” using the asymptotic conditions near the $\mathcal{J}^+$. The scheme consists of two steps [7, 9]:

(i) Propagate the asymptotic generators ξ inward along the null surface Γ intersecting $\mathcal{J}^+$ in Σ^+ by means of geodesic deviation equation

$$\nabla_l^2 \xi + R(\xi, l)l = 0, \quad (11)$$

imposing the following conditions at $\mathcal{J}^+$:

$$\begin{aligned} \lim_{r \rightarrow \infty} l^\mu l^\nu \mathcal{L}_\xi g_{\mu\nu} &= 0, & \lim_{r \rightarrow \infty} r l^\mu m^\nu \mathcal{L}_\xi g_{\mu\nu} &= 0, \\ \lim_{r \rightarrow \infty} l^\mu n^\nu \mathcal{L}_\xi g_{\mu\nu} &= Q(u, \zeta, \bar{\zeta}), \end{aligned}$$

$\mathcal{L}_\xi$ being Lie derivative with respect to ξ .

(ii) Use the commutation relations

$$\begin{aligned} [\xi_a, \xi_b] &= 0, & [\xi_a, \xi_B] &= C_{aB}^b(r, u, \zeta, \bar{\zeta})\xi_b, \\ [\xi_A, \xi_B] &= C_{AB}^D(r, u, \zeta, \bar{\zeta})\xi_D, \end{aligned} \quad (12)$$

and asymptotic expansion of C_{bc}^a

$$C = C_0 + C_1 r^{-1} + C_2 r^{-2} + \dots,$$

r being a canonical parameter and C_0 -s the same as in Eq.(10), evaluate all coefficients of these series. The computation yields [7]:

(a) The generators of translations are given by

$$\xi_a = B_a \Delta^0, \quad (13)$$

where $B_a = l_a(\zeta, \bar{\zeta})/|V(u, \zeta, \bar{\zeta})|$, l_a satisfies $\bar{\partial}_0^2 l_a = 0$ and a “standard edth” $\bar{\partial}_0$ is referred to Bondi frame. There are four independent solutions of this equation.

(b) The generators of boosts and rotations are given by

$$\xi_A = B_A \Delta^0 + C_A \bar{\delta}^0 + \bar{C}_A \delta^0, \quad (14)$$

where $B_A = \Re \tilde{B}_A$, and the complex function $\tilde{B}_A$ satisfies

$$\bar{\partial}^2 \tilde{B}_A - \tilde{B}_A \bar{\lambda}^0 = \frac{\sigma^0}{2} (3\bar{\partial} \bar{C}_A - \bar{\partial} C_A) + \bar{C}_A \bar{\partial} \sigma^0 + C_A \bar{\partial} \sigma^0, \quad (15)$$

σ^0 being the asymptotic shear, λ^0 the news function, and $\bar{\partial} C_A = 0$. There are six independent solutions of this equation which can be written as $C_A = l_A(\zeta, \bar{\zeta})/\bar{V}(u, \zeta, \bar{\zeta})$, where l_A is a solution of the equation $\bar{\partial}_0 l_A = 0$. Further it is convenient to present the function B_A as

$$B_A = \bar{\partial} \eta \bar{C}_A + \frac{u - \eta}{2} \bar{\partial} \bar{C}_A + c.c., \quad (16)$$

(c) The generators of the Poincaré quasigroup obey at $\mathcal{J}^+$ the commutation relations:

$$\begin{aligned} [\xi_a, \xi_b] &= 0, & [\xi_a, \xi_B] &= C_{aB}^b(u, \zeta, \bar{\zeta})\xi_b, \\ [\xi_A, \xi_B] &= C_{AB}^D(u, \zeta, \bar{\zeta})\xi_D, \end{aligned}$$

C_{AB}^D , C_{aB}^i being the *structure functions*, depending on $B_A = \Re \tilde{B}_A$, where $\tilde{B}_A$ is the solution of the equation (15). The last commutation relations mean that the generators of Lorentz quasigroup form a closed algebra.

Note that the same results (a) - (b) can be obtained using instead of (ii) the asymptotic Killing equations $\mathcal{L}_\xi g_{\mu\nu} = 0(1/r^n)$, where the positive integer n depends on the choice of components. This is the usual way emerging the NU (or BMS) group (see, *e.g.*, [24] and references therein). More exactly these equations read

$$\begin{aligned} n^\mu n^\nu \mathcal{L}_\xi g_{\mu\nu} &= O(r), & m^\mu n^\nu \mathcal{L}_\xi g_{\mu\nu} &= O(r), \\ m^\mu m^\nu \mathcal{L}_\xi g_{\mu\nu} &= O(r^0), & m^\mu \bar{m}^\nu \mathcal{L}_\xi g_{\mu\nu} &= O(r^{-2}), \end{aligned} \quad (17)$$

If involve the next orders in powers of r , the system (17) does not have solution, without additional assumptions. It can be understood from the following observation: existence of the solution of the Killing equations for an arbitrary order in powers of r implies that the spacetime has the isometries. Solutions of the geodesic deviation equation exist in an arbitrary spacetime, while the system of the asymptotic Killing equations is compatible with (i), (ii) only for $n \leq 2$ [7, 9]. Note also, that the Jacobi fields play a fundamental role in quasigroup approach to quasilocal conservation quantities [2, 6, 7, 9].

As known a group of isometries can be defined as a group which transforms an arbitrary geodesic to a geodesic one and the Killing vectors satisfy the geodesic deviation equation for any geodesic. In the construction above only *null* geodesics passing inward are transformed to the geodesics under the transformations of the Poincaré quasigroup. Besides, the Poincaré *quasigroup* is appeared only in a *radiative* case. For non-radiating at $\mathcal{J}^+$ systems the generators (13), (14) coincide with these ones obtained by Moreschi [25] via solution of the twistor equation on $\mathcal{J}^+$, and the Poincaré quasigroup is isomorphic to the Poincaré group. It conforms to the well known results on the reduction of the BMS group to the Poincaré group for the asymptotically flat stationary spacetime.

Isometries compatible with gravitational radiation and the Poincaré quasigroup. There are known spacetimes

with double (boost-axial) symmetries which are asymptotically flat and admit radiation [26, 27]. Let us introduce on $\mathcal{J}^+$ the NP basis

$$n|_{\mathcal{J}^+} \equiv \Delta^0 = \partial/\partial u, \\ m|_{\mathcal{J}^+} \equiv \delta^0 = (1/\sqrt{2})(\partial/\partial\theta - (i/\sin\theta)\partial/\partial\varphi).$$

It is easy to show that for the rotational Killing vector (15) is identity. For the boost Killing vector along the axis z Eq.(15) leads to the following constraint on the news function [7, 9]:

$$u \cot\theta \frac{\partial \bar{\lambda}^0}{\partial u} + \frac{\partial \bar{\lambda}^0}{\partial\theta} + 2\bar{\lambda}^0 \cot\theta = 0. \quad (18)$$

This equation is exactly the differential equation for the news function (Eq.(58) in [27]), and hence the quasigroup approach conforms to the known data on the radiative structure of boost-axial spacetime.

Energy-momentum and angular momentum on $\mathcal{J}^+$. As known the Komar integral is not invariant under a change of the choice of the generators of time translations in the equivalence class associated with the given BMS translation. Besides, the resulting energy would not be the monotonically decreasing Bondi energy but the less physical Newman-Unti energy [18]. For an asymptotically flat at future null infinity spacetime the modified “gauge invariant” Komar integral (linkage)

$$L_\xi(\Sigma) = - \lim_{\Sigma_\alpha \rightarrow \mathcal{J}^+} \frac{1}{4\pi} \oint_{\Sigma_\alpha} \left(\xi^{[\alpha;\beta]} + \xi_{;\rho}^\rho l^{[\alpha} n^{\beta]} \right) ds_{\alpha\beta}, \quad (19)$$

where $\{\Sigma_\alpha\}$ being one-parameter family of spheres, was introduced by Tamburino and Winicour [10]. We adopt this as our definition of the conserved quantities on $\mathcal{J}^+$ associated with the generators of the Poincaré quasigroup. The computation leads to the following coordinate independent expression [7, 9]:

$$L_\xi = -(1/4\pi) \Re \oint \left[B(\Psi_2^0 + \sigma^0 \lambda^0 - \bar{\sigma}^2 \bar{\sigma}^0) + 2\bar{C}(\Psi_1^0 + \sigma^0 \bar{\sigma} \bar{\sigma}^0 + (1/2)\bar{\sigma}(\sigma^0 \bar{\sigma}^0)) \right] d\Omega, \quad (\bar{\sigma}C = 0). \quad (20)$$

The integral four-momentum is given by

$$P_i = -(1/4\pi) \Re \oint [B_i(\Psi_2^0 + \sigma^0 \lambda^0 - \bar{\sigma}^2 \bar{\sigma}^0)] d\Omega, \quad (21)$$

where $B_i = l_i/|V|$ and four-vector

$$l = \frac{1}{1+|\zeta|^2} (1+|\zeta|^2, \zeta + \bar{\zeta}, i(\zeta - \bar{\zeta}), |\zeta|^2 - 1).$$

Using the Bianchi identities we compute the loss of energy-momentum

$$\dot{P}_i = -(1/4\pi) \oint B_i |\mathcal{N}|^2 d\Omega, \quad (22)$$

where $\mathcal{N} = \lambda^0 + 2\bar{\sigma}\alpha^0 - 4(\alpha^0)^2$. In the special coordinates ($V = \bar{V}$) the expression (22) coincide with the

expression obtained in [28] and in the Bondi frame we get the standard expression (see, e. g. [29])

$$\dot{P}_i = -(1/4\pi) \oint l_i |\lambda^0|^2 d\Omega. \quad (23)$$

The angular momentum is given by

$$M_A = -(1/4\pi) \Re \oint \bar{l}_A \mathcal{F} d\Omega, \quad (24)$$

where l_A is the solution of $\bar{\sigma}l_A = 0$ and

$$\mathcal{F} = 2\Psi_1^0 + 2\sigma^0 \bar{\sigma} \bar{\sigma}^0 + \bar{\sigma}(\sigma^0 \bar{\sigma}^0) + 3\bar{\sigma}\eta(\Psi_2^0 + \sigma^0 \lambda^0 - \bar{\sigma}^2 \bar{\sigma}^0) + (\eta - u)(\bar{\sigma}\Psi_2^0 + \bar{\sigma}(\sigma^0 \lambda^0) - \bar{\sigma}^3 \bar{\sigma}^0).$$

It yields

$$\dot{M}_A = -(1/4\pi) \Re \oint \bar{l}_A \frac{\partial}{\partial u} \left(\frac{\mathcal{F}}{V|V|^2} \right) V|V|^2 d\Omega. \quad (25)$$

Let Σ_1 and Σ_2 be arbitrary cross-sections at $\mathcal{J}^+$, then the flux of the energy-momentum and angular momentum is given by

$$P_i(\Sigma_2) - P_i(\Sigma_1) = -(1/4\pi) \oint B_i |\mathcal{N}|^2 d\Omega du, \quad (26)$$

$$M_A(\Sigma_2) - M_A(\Sigma_1) = -(1/4\pi) \Re \oint \bar{l}_A \frac{\partial}{\partial u} \left(\frac{\mathcal{F}}{V|V|^2} \right) V|V|^2 d\Omega du, \quad (27)$$

where the integration is performed over the domain $\Omega \subset \mathcal{J}^+$ contained between Σ_1 and Σ_2 .

Geroch and Winicour [30] have given a list of properties which conserved quantities $P(\xi, \Sigma)$ defined at $\mathcal{J}^+$ should have:

- (1) $P(\xi, \Sigma)$ should be linear in the generators of the asymptotic symmetry group.
- (2) $P(\xi, \Sigma)$ should be invariant with respect to the conformal transformations $\tilde{g}_{\mu\nu} = \tilde{\Omega}^2 g_{\mu\nu}$.
- (3) The expression $P(\xi, \Sigma)$ should depends on the geometry of $\mathcal{J}^+$ and behavior of generators in the neighbourhood of $\mathcal{J}^+$.
- (4) $P(\xi, \Sigma)$ should be proportional to the corresponding Komar integral for the exact symmetries and coincide with the Bondi four-momentum when ξ is a BMS translation.
- (5) $P(\xi, \Sigma)$ should be define also for the system with radiation on $\mathcal{J}^+$.
- (6) There should exist a flux integral $\mathcal{I}$ which is linear in ξ and which gives the difference $P(\xi, \Sigma') - P(\xi, \Sigma)$, for Σ' and Σ closed two-surfaces on $\mathcal{J}^+$.
- (7) In Minkowski spacetime $P(\xi, \Sigma)$ should vanish identically.

Our definition of the conserved quantities is free from the supertranslation ambiguity and satisfies all conditions (1) - (7) [7]. In the linearized theory, for the non-contorted two-surfaces, the expression (24) coincide with this one obtained by Bramson [31].

Acknowledgments

I would like to thank N. V. Mitskievich, L. V. Sabinin, T. Zannias, R. D. Sorkin and D. E. Sudarsky for helpfull

discussions and comments. This work was supported by CONACyT, Grant No. 1626P-E.

-
- [1] N. V. Mitskievich, *Physical Fields in General Relativity* (Nauka, Moscow, 1969).
 - [2] N. V. Mitskievich, A. P. Yefremov and A. I. Nesterov, *"Dynamics of Fields in General Relativity"* (Energoatomizdat, Moscow, 1985).
 - [3] D. Bak, Cangemi and R. Jackiw, Phys. Rev. **D 49**, 5173 (1994).
 - [4] A. Komar, Phys. Rev. **113**, 934 (1959).
 - [5] R. Wald, Phys. Rev. **D 48**, R3427 (1993).
 - [6] A. I. Nesterov, *Quasigroups of transformations and conservation laws in General Relativity* (Honorable mention by the Gravity Research Foundation, 1988. Unpublished).
 - [7] A. I. Nesterov, *Methods of Nonassociative Algebra in Physics*. Dr. Sci. Theses (Institute of Physics, Tartu, 1989).
 - [8] A. I. Nesterov. In: *Quasigroups and Nonassociative algebras in Physics* (Insitute of Physics, Tartu, 1990).
 - [9] A. I. Nesterov, *Asymptotic Symmetries and Conservation Laws*. In: *Gravitation and Cosmology* (Moscow, VINITI, 1992).
 - [10] L. A. Tamburino and J. H. Winicour, Phys. Rev. **150**, 1039 (1966).
 - [11] I. A. Batalin, J. Math. Phys. **22**, 1837 (1981).
 - [12] In fact Batalin introduced not a quasigroup but a loop of transformations. The theory of smooth loops generalizing the Lie theory was developed by Sabinin. [13, 14, 15]
 - [13] L. V. Sabinin, In: *Proceedings of Sem. on Vector and Tensor Analysis*, **23** (Moscow Univ., Moscow, 1988).
 - [14] L. V. Sabinin, *Analytic Quasigroups and Geometry* (People's Friendship University, Moscow, 1991).
 - [15] L. V. Sabinin, Russian Math. Survey, **49**, 172 (1994).
 - [16] A. I. Nesterov and L. V. Sabinin, Int. J. Theor. Phys. **36**, 1981 (1997).
 - [17] A. I. Nesterov and L. V. Sabinin, Hadr. J. **20**, 219 (1997).
 - [18] E. T. Newman and T. Unti, J. Math. Phys. **3**, 891 (1962).
 - [19] H. Bondi, A. W. K. Metzner and M. J. G. Van der Berg, Proc Roy. Soc. Lond. **A 269**, 21 (1962).
 - [20] R. K. Sachs, Proc Roy. Soc. Lond. **A 270**, 2851 (1962).
 - [21] R. K. Sachs, Phys. Rev. **128**, 2851 (1962).
 - [22] R. Penrose and W. Rindler *Spinors and space-time* (Cambridge Univ. Press, Cambridge, Vol. I, 1984; Vol. II, 1986).
 - [23] Note, that the last commutation relation can be written as $[\xi_A, \xi_B] = C_{AB}^D \xi_D + C_{AB}^a(u, \zeta, \bar{\zeta}) \xi_a$, where C_{AB}^D are the *structure constants* of the Lorentz group and $C_{AB}^a(u, \zeta, \bar{\zeta})$ the *structure functions* depending on Q .
 - [24] E. T. Newman and K. P. Tod. In: *General Relativity and Gravitation* (Edited by A. Held, Plenum Press, NY, 1980).
 - [25] O. M. Moreschi, Class. Quant. Grav. **3**, 503 (1986).
 - [26] W. Bonnor and N. Z. Swaminarayan, Z. Phys., 240 (1964).
 - [27] J. Bičák and B. G. Schmidt, J. Math. Phys. **25**, 600 (1984).
 - [28] R. W. Lind, J. Messmer and E. T. Newman, J. Math. Phys. **13**, 1884 (1972).
 - [29] J. H. Winicour. In: *General Relativity and Gravitation*, edited by A. Held (Plenum, N.Y. - Lond. 1980).
 - [30] R. Geroch and J. H. Winicour, J. Math. Phys. **22**, 803 (1981).
 - [31] B. Bramson, Proc Roy. Soc. Lond. **A 341**, 463 (1975).